

## **Ocrevus Enrollment Form**

Fax Referral To: 866-832-7180 Phone: 866-852-0202

To better serve your patient and facilitate insurance  • Patient demographics, H&P, and labs • Labs – Antibody testing, BUN/SCr, IgA, IgG  • The front and back of insurance & prescription drug card  • Patient demographics, H&P, and labs • Labs – Antibody testing, BUN/SCr, IgA, IgG  (including subclasses) levels											
PATIENT INFORMATION			PRESCRIBER INFORMATION								
Patient Name:											
Address:City, State, Zip:		City, St	_ City, State, Zip:								
	Cell Phone: Gender:			F <u>ax:</u> NPL#:							
	Phone:			1911#.							
CLINICAL INFORMATION											
DIAGNOSIS: PATIENT EVALUATION:											
G35 Primary Progressive Multiple Sclerosis			Has patient previously received Ocrevus?								
G35 Relapsing Remitting Multiple Sclerosis			If so, date of last infusion with Ocervus: Next dose due:								
			Patient Weight: kg lbs Height: cm in Allergies:								
Required Hepatitis B virus screening results:  Negative Positive (Contraindicated) Date Read:											
PRESCRIPTION INFORMATION											
MEDICATION	DOSE		QUANTITY DISPENSE	REFILLS							
☐ Ocrevus	☐ INITIAL: 300mg IV on day 1 and day 15			Dispense day 1 and day 15	0						
☐ Maintenance: 600mg IV every 6 months starting 6 months after day 1 dose			s after day 1 dose	1 cycle	ONE						
To be infused peripherally, unless otherwise indicated.  PORT PICC  • 0.9% NaCl: 1-5ml intravenous (peripheral line) or 10ml intravenous (central line) before/after infusion, or as needed for line patency/SASH.  • Heparin 10 units/ml: 5ml intravenous (peripheral line) as needed for final flush.  • Heparin 100 units/ml: 5ml intravenous (central line) as needed for final flush.											
HYDRATION:											
						PREMEDICATIONS: to be given 30 minutes prior to infusion (PLEASE STRIKE THROUGH IF NOT REQUIRED)					
						Diphenhydramine 25mg-50mg by mouth Acetaminophen 325mg-650mg by mouth Methylprednisolone 100mg slow IV push					
<ul><li>Diphenhydrar</li><li>Acetaminophe</li></ul>	o be used as needed/requested by patie nine: May repeat premedication dose e en: May repeat premedication dose even 6/Prilocaine 2.5% topical (may dispense	every 4-6 hours as needed. Adult r ry 4-6 hours as needed. Adult ma	nax: 100mg/day. x: 3000mg/day.								
	PROTOCOL: Pharmacy to provide the										
<ul> <li>Diphenhydramine 25mg- 50mg PO as needed for mild-moderate reaction. qty: 2</li> <li>Diphenhydramine 25mg- 50mg via slow IV push as needed for moderate-severe</li> <li>Epinephrine 0.3mg (patient's weighing ≥ 30kg/66lbs)</li> <li>Epinephrine 0.15mg (patient's weighing between 15-30kg/33-66lbs)</li> </ul>											
reaction. qty: one 50mg/ml vial  • Sodium Chloride 0.9% 500ml bag IV PRN for anaphylactic reaction. qty: 500ml  Administer intramuscularly PRN severe allergic reaction.  Call 911. May repeat x 1.											
DELIVERY MET	HOD: To be infused via gravity infusion maintenance infusions to be titrated	on, unless otherwise indicated	☐ MD prefers Infusion								
NURSING: Pharmacy to coordinate home health nursing visit as necessary:											
ADDITIONAL:											
By signing this form and using this pharmacy's services, you are authorizing this pharmacy to serve as your prior authorization designated agent in dealing with prescription and medical insurance companies. 🔲 Dispense as Written											

\_ Date: \_\_\_

Prescriber Signature: \_\_

☐ Product Substitution Permitted